REFERENCES

Botanical Name: Microstegium vimineum

Common Name: Japanese stiltgrass, Nepalese browntop

Family Name: Poaceae

- 1. USDA Plants database, plant profiles: http://plants.usda.gov/java/nameSearch. Accessed 8-13-12
- 2. EDDMapS (2012) Early Detection & Distribution Mapping System. The University of Georgia Center for Invasive Species and Ecosystem Health. Available online at http://www.eddmaps.org/; last accessed June 27, 2012.
- 3. Indiana's "Most Wanted" Invasive Plant Pests: Indian Cooperative Agricultural Pest Survey (CAPS) Program: http://extension.entm.purdue.edu/caps/browsePest.html. Accessed8-13-12.
- 4. Kentucky Exotic Pest Plant Council: http://www.se-eppc.org/ky/list.htm. Accessed 8-13-12.
- 5. Michigan State University Extension; The Michigan Natural Features Inventory (MNFI) has partnered with MISIN to provide the information in this fact sheet. Original content was taken with permission from the MNFI field guide entitled: A Field Identification Guide to Invasive Plants in Michigan's Natural Communities (PDF).: http://mnfi.anr.msu.edu/education/factsheets.cfm. Accessed on8-13-12
- 6. Pennsylvania Dept. Of Conservation and Natural Resources: Invasive Plants in Pennsylvania: http://www.dcnr.state.pa.us/ucmprd2/groups/public/documents/document/dcnr_002477.pdf. Accessed 8-13-12.
- 7. Adams, S.N. and K.A.M. Engelhardt(2009) Diversity declines in microstegium vimineum patches. Biological Conservation 142: 1003-1010.
- 8. Huebner, C.D. (2011) Seed mass, viability, and germination of Japanese stiltgrass (Microstegium vimineum) under variable light and moisture conditions". Invasive Plant Science and Management 4(3): 274.
- 9. Ward, J.S. and Mervosh, T.L. (2012). Nonchemical and herbicide treatments for management of Japanese stiltgrass (Microstegium vimineum). Invasive Plant Science and Management 5(1): 9.
- 10. Fryer, J.L. (2011) Microstegium vimineum. In: Fire Effects Information System, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). Available: http://www.fs.fed.us/database/feis/ [2012, August 13].
- 11. Abrams, M.D., and S.E. Johnson (2012) Long-term impacts of deer exclosures on mixed-oak forest composition at the Valley Forge National Historical Park, Pennsylvania, USA. Journal of the Torrey Botanical Society 139(2): 167–180.
- 12. Adams, S.N., and K.A.M. Engelhardt (2009) Diversity declines in Microstegium vimineum (Japanese stiltgrass) patches. Biological Conservation 142: 1003–1010.
- 13. Anderson, D.P., M.G. Turner, S.M. Pearson, T.P. Albright, R.K. Peet, and A. Wieben (2013) Predicting Microstegium vimineum invasion in natural plant communities of the southern Blue Ridge Mountains, USA. Biol Invasions 15: 1217–1230.
- 14. Aronson, M.F.J., and S.N. Handel (2011) Deer and Invasive Plant Species Suppress Forest Herbaceous Communities and Canopy Tree Regeneration. Natural Areas Journal 31(4): 400-407.
- 15. Bauer, J.T. (2012) Invasive species: "back-seat drivers" of ecosystem change? Biol Invasions 14: 1295-1304.

- 16. Bauer, J.T. and S.L. Flory (2011) Suppression of the Woodland Herb Senna hebecarpa by the Invasive Grass Microstegium vimineum. The American Midland Naturalist 165(1): 105-115.
- 17. Beasley,R.R. and B.C. McCarthy (2011) Effects of Microstegium vimineum (Trin.) A. Camus (Japanese Stiltgrass) on Native Hardwood Survival and Growth: Implications for Restoration. Natural Areas Journal 31(3): 246-255.
- 18. Bradford, M.A., M.S. Strickland, J.L. DeVore, and J.C. Maerz (2012) Root carbon flow from an invasive plant to belowground foodwebs. Plant Soil 359: 233–244.
- 19. Cheplick, G.P. and J. Fox (2011) Density-dependent growth and reproduction of Microstegium vimineum in contrasting light environments. The Journal of the Torrey Botanical Society 138(1): 62-72.
- 20. Cheplick, G.P. (2010) Limits to local spatial spread in a highly invasive annual grass (Microstegium vimineum). Biol Invasions 12: 1759–1771.
- 21. Cipollini, K.A. and D. Cipollini (2011) Habitat Assessment and Conservation Status of Endangered Northeastern Bulrush. Northeastern Naturalist 18(3): 275-291.
- 22. Corbett, B.F., and J.A. Morrison (2012) The Allelopathic Potentials of the Non-Native Invasive Plant Microstegium vimineum and the Native Ageratina altissima: Two Dominant Species of the Eastern Forest Herb Layer. Northeastern Naturalist 19(2): 297-312.
- 23.Emery, S.M., J. Uwimbabazia, and S.L. Flory (2011) Fire intensity effects on seed germination of native and invasive Eastern deciduous forest understory plants. Forest Ecology and Management 261: 1401–1408.
- 24.Eschtruth, A.K., and J.J. Battles (2011) The importance of quantifying propagule pressure to understand invasion: an examination of riparian forest invasibility. Ecology 92(6): 1314–1322.
- 25. Flory, S.L., and K. Clay (2010) Non-native grass invasion suppresses forest succession. Oecologia 164: 1029–1038.
- 26. Fraterrigo, J.F., M.S. Strickland, A.D. Keiser, and M.A. Bradford (2011) Nitrogen uptake and preference in a forest understory following invasion by an exotic grass. Oecologia 167: 781–791.
- 27. Hayes, S.J. and E.J. Holzmueller (2012) Relationship between Invasive Plant Species and Forest Fauna in Eastern North America. Forests 3: 840-852.
- 28. Nagy, C., S. Aschen, R. Christie, and M. Weckel (2011) Japanese stilt grass (Microstegium vimineum), a nonnative invasive grass, provides alternative habitat for native frogs in a suburban forest. Urban Habitats. Available online: http://www.urbanhabitats.org/v06n01/japanesestiltgrass_full.html Accessed 12/4/2103.
- 29. Huebner, C.D. (2010) Establishment of an invasive grass in closed-canopy deciduous forests across local and regional environmental gradients. Biol Invasions 12: 2069–2080.
- 30. Kleczewski, N.M., S.L. Flory, and K. Clay (2012) Variation in Pathogenicity and Host Range of Bipolaris sp. Causing Leaf Blight Disease on the Invasive Grass Microstegium vimineum. Weed Science 60(3): 486-493.
- 31. Kramer, T.D., R.J. Warren, Y. Tang, and M.A. Bradford (2012) Grass Invasions Across a Regional Gradient are Associated with Declines in Belowground Carbon Pools. Ecosystems 15: 1271–1282.
- 32. Kuebbing, S., M.A. Rodriguez-Cabal, D. Fowler, L. Breza, J.A. Schweitzer and J.K. Bailey (2012) Resource availability and plant diversity explain patterns of invasion of an exotic grass. Journal of Plant Ecology doi: 10.1093/jpe/rts018
- 33. Lee, M.R., S.L. Flory, and R.P. Phillips (2012) Positive feedbacks to growth of an invasive grass through alteration of nitrogen cycling. Oecologia 170: 457–465.
- 34. Novy, A., S.L. Flory, and J.M. Hartman (2013) Evidence for rapid evolution of phenology in an invasive grass. J. Evol. Biol. 26: 443–450.
- 35. Schramm, J.W., and J.G. Ehrenfeld (2012) Patterns of patch colonization and explansion in the non-native annual grass Microstegium vimineum (Poaceae). Rhodora 114(957): 1–20.
- 36. Simao, M.C.M., S.L. Flory and J.A. Rudgers (2010) Experimental plant invasion reduces arthropod abundance and richness across multiple trophic levels. Oikos 119: 1553–1562.

- 37.Strickland, M.S., J.L. DeVore, J.C. Maerz, and M.A. Bradford (2010) Loss of faster-cycling soil carbon pools following grass invasion across multiple forest sites. Soil Biology & Biochemistry 43: 452-454.
- 38. Tang, Y., R.J. Warren, T.D. Kramer, and M.A. Bradford (2012) Plant invasion impacts on arthropod abundance, diversity and feeding consistent across environmental and geographic gradients. Biol Invasions 14: 2625–2637.
- 39. Tekiela, D.R. and J.N. Barney (2013) Quantifying Microstegium vimineum Seed Movement by Non-Riparian Water Dispersal Using an Ultraviolet-Marking Based Recapture Method. PLOS ONE 8(9): e63811.
- 40. Ward, J.S. and T.L. Mervosh (2012) Nonchemical and Herbicide Treatments for Management of Japanese Stiltgrass (Microstegium vimineum). Invasive Plant Science and Management 5:9-
- 41. Warren, R.J., J.P. Wright, and M.A. Bradford (2011) The putative niche requirements and landscape dynamics of Microstegium vimineum: an invasive Asian grass. Biol Invasions 13: 471–483.
- 42. Warren, R.J., V. Bahn, and M.A. Bradford (2012) The interaction between propagule pressure, habitat suitability and density-dependent reproduction in species invasion. Oikos 121: 874–881.
- 43. Warren, R.J., V. Bahn, and M.A. Bradford (2013) Decoupling litter barrier and soil moisture influences on the establishment of an invasive grass. Plant Soil 367: 339–346.